

2008

Length-Weight Relationships of Select Common Nearshore Southern California Marine Fishes

Eric F. Miller

D. S. Beck

Wayne Dossett

Follow this and additional works at: <https://scholar.oxy.edu/scas>

Part of the [Ecology and Evolutionary Biology Commons](#), and the [Marine Biology Commons](#)

Recommended Citation

Miller, Eric F.; Beck, D. S.; and Dossett, Wayne (2008) "Length-Weight Relationships of Select Common Nearshore Southern California Marine Fishes," *Bulletin of the Southern California Academy of Sciences*: Vol. 107: Iss. 3.
Available at: <https://scholar.oxy.edu/scas/vol107/iss3/3>

This Article is brought to you for free and open access by OxyScholar. It has been accepted for inclusion in Bulletin of the Southern California Academy of Sciences by an authorized editor of OxyScholar. For more information, please contact cdla@oxy.edu.

Length-Weight Relationships of Select Common Nearshore Southern California Marine Fishes

Eric F. Miller,¹ D. Shane Beck, and Wayne Dossett

MBC Applied Environmental Sciences, 3000 Red Hill Ave., Costa Mesa, CA 92626

Growth dynamics of marine fish is often critical for fisheries analysis. Frequently only lengths are measured during ecological surveys, occasionally with aggregate weights by species (Allen and Herbinson 1990; Pondella and Allen 2000; Allen et al. 2002). This information often allows for estimation of age-class composition through length frequency analysis, but can mask the importance of individual fish weights to the overall community structure, such as the inclusion of one, or a few, large individuals among a predominantly small catch. Over time, authors have reported length-to-weight relationships for specific species in the course of specific life history investigations, but these studies are rarely conducted on forage species. Cailliet et al. (2000) compiled many of the relationships that were available at the time, but numerous species still lack such basal information. Their database, however, contained 124 species, many of which did not have length-weight relationships available. Love et al. (2002) also compiled additional information on several rockfish species (*Sebastes* spp) and *Sebastolobus alascamus*.

Since 1979, impingement surveys at coastal generating stations routinely recorded the length of nearly all impinged fish during a given survey. Most of these surveys, however, recorded the aggregate weight by species, a protocol consistent with most ecological studies in southern California. Specific, focused studies were occasionally undertaken during which the individual length and weight of a subset of individuals were recorded. Moreover, instances when only a single individual of a specific species was impinged, a defacto length-weight relationship data point was recorded. During these studies the appropriate length; standard (SL), total (TL), or disc width (DW) was measured to the nearest millimeter (mm) and weight recorded to the nearest gram (g).

Data from impingement records 2001–2006, recorded at generating stations from San Diego County to Los Angeles County, California were reviewed to generate species-specific length-weight relationships (Tables 1 and 2). A total of 59 species were identified with length and weight recorded for greater than 10 individuals. These included both 54 ray-finned fishes (Class Actinopterygii) and 5 elasmobranch species (Subclass Elasmobranchii). All were common to the Southern California Bight (Miller and Lea 1972; Love et al. 2005). Forty-two of these species were included in the Cailliet et al. (2000) species list, but only 27 had a length-weight relationship listed.

The length-weight relationships of fishes typically fit the non-linear equation $W = aL^b$, where W = weight (g), L = length (mm), and a and b are derived constants. A best fit line was plotted for each distribution using MS Excel. The determination of the best fit was based on the R^2 -value. Fifty-six species were best described by the traditional non-linear equation $W = aL^b$. Two species, *Anchoa compressa* and *A. delicatissima* were best described by a linear relationship and one species, *Seriphus politus*, was best described by an exponential function. The minimum and maximum length (mm) recorded for each

¹ Corresponding Author: emiller@mbcnet.net

Table 1. Length (mm)-weight (g) relationship, tightness of fit to the equation (R^2), sample size, and minimum and maximum length recorded for 54 fish species (Class Actinopterygii) collected during impingement sampling from Los Angeles County to San Diego County, California from 2001–2006, mainly 2006.

Species	Equation	R^2	N	Min. Len. (mm)	Max. Len. (mm)
<i>Acanthogobius flavimanus</i>	Wt = 2E-05SL ^{2.8272}	0.84	67	70	183
<i>Anchoa compressa</i>	Wt = 0.3SL-0.0164	0.78	187	62	126
<i>Anchoa delicatissima</i>	Wt = 0.2SL - 0.0123	0.76	119	40	129
<i>Anisotremus davidsonii</i>	Wt = 2E-05SL ^{3.0457}	0.99	45	41	327
<i>Atherinops affinis</i>	Wt = 2E-06SL ^{3.3046}	0.96	1499	22	286
<i>Atherinopsis californiensis</i>	Wt = 2E-055SL ^{2.9051}	0.89	668	48	342
<i>Atractoscion nobilis</i>	Wt = 4E-05SL ^{2.7991}	0.98	65	64	420
<i>Brachyistius frenatus</i>	Wt = 5E-06SL ^{3.3596}	0.93	22	70	128
<i>Chilara taylori</i>	Wt = 1E-06SL ^{3.2744}	0.92	18	140	226
<i>Chromis punctipinnis</i>	Wt = 2E-05SL ^{3.1244}	0.97	133	28	238
<i>Citharichthys stigmaeus</i>	Wt = 2E-05SL ^{3.0021}	0.87	165	22	116
<i>Cymatogaster aggregata</i>	Wt = 2E-05SL ^{3.0583}	0.95	824	29	161
<i>Embiotoca jacksoni</i>	Wt = 8E-06SL ^{3.3244}	0.98	413	47	256
<i>Fundulus parvipinnis</i>	Wt = 8E-05SL ^{2.6197}	0.87	14	48	70
<i>Genyonemus lineatus</i>	Wt = 1E-05SL ^{3.1249}	0.98	389	28	252
<i>Gibbonsia elegans</i>	Wt = 6E-06SL ^{3.171}	0.92	32	48	160
<i>Gillichthys mirabilis</i>	Wt = 1E-04SL ^{2.4415}	0.87	112	25	154
<i>Girella nigricans</i>	Wt = 2E-05SL ^{3.124}	0.98	21	75	332
<i>Halichoeres semicinctus</i>	Wt = 5E-06SL ^{3.2776}	0.95	33	136	286
<i>Heterostichus rostratus</i>	Wt = 7E-06SL ^{3.0407}	0.95	131	33	325
<i>Hyperprosopon argenteum</i>	Wt = 1E-05SL ^{3.2295}	0.97	274	42	168
<i>Hypsoblennius gilberti</i>	Wt = 6E-05SL ^{2.7702}	0.88	87	31	113
<i>Hypsurus caryi</i>	Wt = 2E-05SL ^{3.051}	0.99	27	50	186
<i>Leptocottus armatus</i>	Wt = 2E-05SL ^{2.9422}	0.94	326	32	185
<i>Leuresthes tenuis</i>	Wt = 7E-06SL ^{3.043}	0.95	125	31	165
<i>Medialuna californiensis</i>	Wt = 3E-05SL ^{3.0106}	0.97	12	145	224
<i>Menticirrhus undulatus</i>	Wt = 2E-05SL ^{2.9383}	0.99	72	56	375
<i>Ophichthus zophochir</i>	Wt = 1E-12SL ^{5.1724}	0.91	11	419	672
<i>Ophidion scrippsae</i>	Wt = 4E-08SL ^{3.9624}	0.96	14	130	249
<i>Oxyjulis californica</i>	Wt = 2E-05SL ^{2.939}	0.92	35	82	180
<i>Paralabrax clathratus</i>	Wt = 4E-05SL ^{2.9184}	0.98	242	25	402
<i>Paralabrax nebulifer</i>	Wt = 3E-05SL ^{2.9802}	0.99	150	27	435
<i>Paralichthys californicus</i>	Wt = 8E-05SL ^{2.6813}	0.88	53	41	508
<i>Peprilus simillimus</i>	Wt = 1E-05SL ^{3.1729}	0.90	160	45	175
<i>Phanerodon furcatus</i>	Wt = 2E-05SL ^{3.053}	0.97	150	33	200
<i>Pleuronichthys guttulus</i>	Wt = 4E-05SL ^{2.9106}	0.99	61	27	375
<i>Pleuronichthys ritteri</i>	Wt = 2E-05SL ^{3.083}	0.97	185	20	245
<i>Pleuronichthys verticalis</i>	Wt = 3E-05SL ^{2.9387}	0.98	40	54	294
<i>Porichthys myriaster</i>	Wt = 1E-05SL ^{2.9407}	0.97	230	19	476
<i>Porichthys notatus</i>	Wt = 8E-06SL ^{3.0692}	0.98	25	37	266
<i>Rhacochilus toxotes</i>	Wt = 2E-05SL ^{3.0787}	0.99	98	68	281
<i>Rhacochilus vacca</i>	Wt = 4E-05SL ^{2.9363}	0.97	69	54	276
<i>Scomber japonicus</i>	Wt = 8E-06SL ^{3.0902}	0.85	272	74	363
<i>Scorpaena guttata</i>	Wt = 4E-05SL ^{2.981}	0.98	145	29	293
<i>Scorpaenichthys marmoratus</i>	Wt = 3E-05SL ^{2.9904}	0.97	37	79	285
<i>Sebastes auriculatus</i>	Wt = 2E-05SL ^{3.0945}	0.95	51	110	316
<i>Sebastes miniatus</i>	Wt = 5E-05SL ^{2.9095}	0.99	12	39	290

Table 1. Continued.

Species	Equation	R ²	N	Min. Len. (mm)	Max. Len. (mm)
<i>Seriphus politus</i>	Wt = 0.3e ^{0.0373SL}	0.93	3073	11	193
<i>Sphyraena argentea</i>	Wt = 5E-06SL ^{3.0316}	0.97	31	45	406
<i>Symphurus atricaudus</i>	Wt = 8E-04SL ^{2.0957}	0.91	18	22	179
<i>Synodus lucioceps</i>	Wt = 4E-06SL ^{3.1072}	0.87	85	26	221
<i>Trachurus symmetricus</i>	Wt = 7E-06SL ^{3.1246}	0.91	159	60	241
<i>Umbrina roncadore</i>	Wt = 3E-058SL ^{2.9233}	0.99	73	72	301
<i>Xenistius californiensis</i>	Wt = 2E-05SL ^{3.0349}	0.94	265	29	173

Table 2. Length (mm)-weight (g) relationships of fit to the equation (R²), sample size, and minimum and maximum length recorded for five elasmobranch (Subclass Elasmobranchii) species collected during impingement sampling from Los Angeles County to San Diego County, California from 2001–2006, mainly 2006.

Species	Equation	R ²	N	Min. Len. (mm)	Max. Len. (mm)
<i>Heterodontus francisci</i>	Wt = 9E-06TL ^{2.9948}	0.98	19	112	750
<i>Myliobatis californica</i>	Wt = 1E-05DW ^{3.0436}	0.95	446	167	885
<i>Platyrrhinoidis triseriata</i>	Wt = 7E-06TL ^{2.9774}	0.98	102	115	670
<i>Torpedo californica</i>	Wt = 7E-05TL ^{2.7748}	0.90	50	185	932
<i>Urobatis halleri</i>	Wt = 3E-05DW ^{3.1312}	0.88	960	68	366

species was included, as some of these relationships may not fully encapsulate the total available size range commonly occurring in the Southern California Bight. These relationships were not compared to previous published results, but rather presented purely based on the available data recorded during impingement surveys so as to represent a recent assessment of length-weight relationships of common marine fish.

Acknowledgements

We would like to thank R. Moore and M. Love, whose questions and support spurred the compilation of this data. This work could not have been possible without the support of the owners and operators of the coastal power plants who funded the impingement surveys: Los Angeles Department of Water and Power, AES Redondo Beach, AES Alamitos, AES Huntington Beach, El Segundo Power, and Southern California Edison. We would like to thank the various staff members of MBC Applied Environmental Sciences from 2001 to 2007, who conducted these surveys during all hours. This manuscript benefited from the comments of D. Guthrie and three anonymous reviewers.

Literature Cited

- Allen, L.G., A.M. Findlay, and C.M. Phalen. 2002. Structure and standing stock of the fish assemblages of San Diego Bay, California from 1994–1999. *Bull. South. Calif. Acad. Sci.*, 101:49–85.
- Allen, M.J. and K.T. Herbinson. 1990. Settlement of juvenile California halibut, *Paralichthys californicus*, along the coasts of Los Angeles, Orange, and San Diego Counties in 1989. *Cal. Coop. Oceanic Fish. Invest. Rep.*, 31:84–96.

- Cailliet, G.M., E.J. Burton, J.M. Cope, L.A. Kerr, R.J. Larson, R.N. Lea, D. Van Tresca, and E. Knaggs. 2000. Biological characteristics of nearshore fishes of California: a review of existing knowledge and proposed additional studies. <http://www.dfg.ca.gov/marine/lifehistories/index.asp>
- Love, M.S., C.W. Mecklenburg, T.A. Mecklenburg, and L.K. Thorsteinson. 2005. Resource Inventory of Marine and Estuarine Fishes of the West Coast and Alaska: A Checklist of North Pacific and Arctic Ocean Species from Baja California to the Alaska-Yukon Border. U. S. Department of the Interior, U. S. Geological Survey, Biological Resources Division, Seattle, Washington, 98104, OCS Study MMS 2005-030 and USGS/NBII 2005-001.
- , M. Yoklavich, and L. Thorsteinson. 2002. The rockfishes of the northeast Pacific. UC Press, Los Angeles, California.
- Miller, D.J. and R.N. Lea. 1972. Guide to the coastal marine fishes of California. Calif. Dep. Fish Game, Fish Bull. 157, 249 pp.
- Pondella, D.J. II and L.G. Allen. 2000. The nearshore fish assemblage of Santa Catalina Island. *In*: The Proceedings of the Fifth California Islands Symposium. (D.R. Browne, K.L. Mitchell, and H.W. Chaney, eds.) Santa Barbara Museum of Natural History, Santa Barbara, California. Pp. 394–400.
- Accepted for publication 22 May 2008.