

9-26-2014

## Psoroptes Mites and Mule Deer (*Odocoileus hemionus*): Additional Notes from the San Bernardino Mountains, California

Vernon C. Bleich

*Idaho State University*, [vcbleich@gmail.com](mailto:vcbleich@gmail.com)

Follow this and additional works at: <http://scholar.oxy.edu/scas>



Part of the [Life Sciences Commons](#)

---

### Recommended Citation

Bleich, Vernon C. (2014) "Psoroptes Mites and Mule Deer (*Odocoileus hemionus*): Additional Notes from the San Bernardino Mountains, California," *Bulletin of the Southern California Academy of Sciences*: Vol. 113: Iss. 2.  
Available at: <http://scholar.oxy.edu/scas/vol113/iss2/4>

This Research Note is brought to you for free and open access by OxyScholar. It has been accepted for inclusion in Bulletin of the Southern California Academy of Sciences by an authorized administrator of OxyScholar. For more information, please contact [cdla@oxy.edu](mailto:cdla@oxy.edu).

---

# Psoroptes Mites and Mule Deer (*Odocoileus hemionus*): Additional Notes from the San Bernardino Mountains, California

## **Cover Page Footnote**

We thank B. Adams, B. Clark, A. Hunter, R. Teagle and T. Glenner — as well as numerous other CDFW personnel — and S. deJesus (Landells Aviation) for help capturing and handling study animals; T. Evans (CDFW) for conducting telemetry flights; and K. Jones and L. Konde (CDFW) for careful record-keeping and archiving of relevant data. We thank D. Shumway and D. Rib (Mitsubishi Cement Corporation), and D. Anderson (Specialty Minerals Corporation) for assistance with logistics and arranging for access to private property. Financial support for this research and for manuscript preparation was provided by the Mitsubishi Cement Corporation, CDFW, Granite Bay Chapter of Safari Club International, and the Eastern Sierra Center for Applied Population Ecology. This is Professional Paper 100 from the Eastern Sierra Center for Applied Population Ecology.

***Psoroptes* Mites and Mule Deer (*Odocoileus hemionus*):  
Additional Notes from the San Bernardino Mountains, California**

Vernon C. Bleich,<sup>1,4</sup> Jeffrey T. Villepique,<sup>2</sup> James H. Davis,<sup>3</sup> Steven G. Torres,<sup>3</sup>  
and Ben J. Gonzales<sup>3</sup>

<sup>1</sup>*California Department of Fish and Wildlife, Sierra Nevada Bighorn Sheep Recovery  
Program, 407 W. Line St., Bishop, CA 93514 USA*

<sup>2</sup>*California Department of Fish and Wildlife, P.O. Box 3222, Big Bear City,  
CA 92314 USA*

<sup>3</sup>*California Department of Fish and Wildlife, Wildlife Investigations Laboratory,  
1701 Nimbus Rd., Rancho Cordova, CA 95670 USA*

---

Psoroptic scabies is caused by a mite (*Psoroptes* sp., Asghar et al., 2011; but, see Zahler et al. 2000) and affects bighorn sheep (*Ovis canadensis*) and cattle in the western United States (Singer et al. 1997). Similar to other diseases shared by native and domestic animals, psoroptic scabies has implications for the animal production industry and the health and conservation of wild ungulates (Roug et al. 2012). *Psoroptes* mites previously have been detected on mule deer (*Odocoileus hemionus*) sympatric with a population of desert bighorn sheep in New Mexico (Lange et al. 1980; Hoban 1990; Logan and Sweanor 2001) that was impacted substantially by that ectoparasite (Boyce and Brown 1991). Mite infestations among desert bighorn sheep in California are widespread and commonplace (Mazet et al. 1992; Clark et al. 1993), yet do not appear to result in severe clinical disease (Mazet et al. 1992).

Several of the populations of bighorn sheep examined by other investigators (Mazet et al. 1992; Clark et al. 1993) were sympatric with mule deer, as was the situation in the San Bernardino Mountains, California (Schaefer et al. 2000). Singer et al. (1997) investigated the prevalence of *Psoroptes* infestation among those species and domestic cattle in the southeastern part of that mountain range. Mule deer and bighorn sheep also were sympatric on the north-facing slope of the San Bernardino Mountains (Cushenbury Canyon, Bleich et al. 2009; Abella et al. 2011), and often share chaparral-dominated habitats in other parts of the transverse ranges and in the peninsular ranges of California (Bleich and Holl 1982). Psoroptic scabies was present in bighorn sheep captured in the San Bernardino Mountains at Cushenbury Canyon (Clark et al. 1988) and near Mt. San Gorgonio (Singer et al. 1997).

Singer et al. (1997) reported an absence of *Psoroptes* infestation among 15 mule deer captured near Mt. San Gorgonio. The small number of deer examined, however, limits the ability to make a conclusion based on negative detection of a pathogen or other anomaly (Hanley and Lippman-Hand 1983; Dougherty and McInerney 2009; Ho 2009). Singer et al. (1997) recognized a need to replicate results and recommended further investigation of pathogen-sharing among sympatric populations of mule deer and bighorn sheep. Therefore, we sampled additional mule deer from the San Bernardino Mountains to further explore the question of whether *Psoroptes* mites infest mule deer

---

<sup>4</sup> Present address: Department of Biological Sciences, Idaho State University, Pocatello, ID 83209 USA  
Corresponding author: vern.bleich@wildlife.ca.gov

Table 1. Sample size and associated level of confidence that the prevalence ( $P$ ) of *Psoroptes* mites was  $\leq 10\%$  in two proximate populations of mule deer, each numbering an estimated 200 individuals and sympatric with bighorn sheep in the San Bernardino Mountains, San Bernardino County, California, 1991–2004.

Source	Date	Approximate study centroid of area <sup>a</sup>	$N$	Level of confidence $P^b \leq 0.10$
Singer et al. (1997)	1991–94	34°04'N, 116°39'W	15	81%
This study	2002	34°21'N, 116°55'W	16	83%
This study	2004	34°21'N, 116°55'W	18	86%

<sup>a</sup> Latitude and longitude, WGS84 datum.

<sup>b</sup> Probability of exposure estimated according to the method of Wehausen (1987).

that are sympatric with bighorn sheep in that range, as originally posited by Singer et al. (1997).

We captured mule deer in the vicinity of Cushenbury Canyon during 2002 and 2004. We used a net-gun fired from a helicopter (Jessup et al. 1988); deer were physically restrained, and some individuals were transported by helicopter to a central base for processing (Monteith et al. 2011; Monteith et al. 2013), but those caught  $>10$  km from that base were processed in the field. Each individual received prophylactic injections of penicillin (Sterile Penicillin Procaine G; Aspen Veterinary Resources, Liberty, MO, USA), Vitamin E (Vital-E®; Schering-Plough Animal Health Corporation, Summit, NJ, USA), and Vitamin E and Selenium (MU-SE®; Schering-Plough Animal Health Corporation, Summit, NJ, USA), and each was fitted with a VHF telemetry collar prior to release. Capture and handling procedures followed protocols developed by the California Department of Fish and Wildlife (CDFW) (Jessup et al. 1986) and were consistent with guidelines for acceptable field methods adopted by the American Society of Mammalogists (Animal Care and Use Committee 1998).

We conducted a thorough physical examination of each deer in an effort to detect clinical evidence of mite infestation, paying careful attention to the ears as described by Welsh and Bunch (1983). We searched both ears of each individual for evidence of dermatitis in the form of exfoliated epidermis, crusted serous exudate, or blockage of the meatus with hard, waxy material that would be consistent with clinical evidence of *Psoroptes* infestation (Welsh and Bunch 1983). We also collected deep ear swabs (Singer et al. 1997), which were examined microscopically for presence of *Psoroptes* mites. None of the mule deer captured in 2002 ( $n = 16$ ) or 2004 ( $n = 18$ ) revealed clinical evidence of *Psoroptes* infestation. Further, deep ear swabs collected from mule deer captured during September 2002 and later examined microscopically ( $n = 5$ ) were negative for mites (K. R. Jones, CDFW, *in litt*, 14 March 2013).

It will be possible to establish an unambiguous absence of *Psoroptes* infestation in mule deer in the San Bernardino Mountains only if every individual can be sampled (Wehausen 1987). Nevertheless, using the method developed by Wehausen (1987), it can be stated with a high degree of confidence ( $\bar{x} = 83.33\% \pm 4.023$  [95% CI]; Table 1) that the infestation rate during each sampling period was  $\leq 10\%$ . Moreover, assuming that the infestation rates in each of the 3 sampling periods were independent, it is highly likely ( $P = 0.9945$ ;  $1 - \{1 - 0.81\} \times \{1 - 0.83\} \times \{1 - 0.87\}$ ) that the infestation rate among the populations of mule deer examined was  $\leq 10\%$ .

Our failure to detect clinical evidence of psoroptic scabies or the presence of *Psoroptes* mites is consistent with the results of Singer et al. (1997), who detected no evidence of *Psoroptes* infestation among the mule deer they examined. Given the similarity of the level of confidence for each sampling effort, it is entirely plausible that no deer in the geographic areas investigated by us or Singer et al. (1997) were infested with mites. Our results provide additional evidence, as emphasized by Singer et al. (1997), that mule deer are an unlikely source of *Psoroptes* mites that infest bighorn sheep inhabiting the San Bernardino Mountains. Nevertheless, we cannot rule out other species of wildlife as potential sources of infestation of bighorn sheep by *Psoroptes* mites.

We thank B. Adams, B. Clark, A. Hunter, R. Teagle and T. Glenner — as well as numerous other CDFW personnel — and S. deJesus (Landells Aviation) for help capturing and handling study animals; T. Evans (CDFW) for conducting telemetry flights; and K. Jones and L. Konde (CDFW) for careful record-keeping and archiving of relevant data. We thank D. Shumway and D. Rib (Mitsubishi Cement Corporation), and D. Anderson (Specialty Minerals Corporation) for assistance with logistics and arranging for access to private property. Financial support for this research and for manuscript preparation was provided by the Mitsubishi Cement Corporation, CDFW, Granite Bay Chapter of Safari Club International, and the Eastern Sierra Center for Applied Population Ecology. This is Professional Paper 100 from the Eastern Sierra Center for Applied Population Ecology.

#### Literature Cited

- Abella, R., V.C. Bleich, R.A. Botta, B.J. Gonzales, T.R. Stephenson, S.G. Torres, and J.D. Wehausen. 2011. Status of bighorn sheep in California — 2011. *Desert Bighorn Council Transactions*, 51: 54–68.
- Asghar, A., O. Hassani, A. Alsadi, H. Feda, and S. Faithi. 2011. Prevalence of scabies diagnosed in sheep and goats during Hajj season in Makkah. *Journal of Agricultural and Veterinary Sciences*, 4: 37–43.
- Animal Care and Use Committee. 1998. Guidelines for the capture, handling, and care of mammals as approved by the American Society of Mammalogists. *Journal of Mammalogy*, 79:1416–1431.
- Bleich, V.C. and S.A. Holl. 1982. Management of chaparral habitat for mule deer and mountain sheep in southern California. Pp. 247–254 in C.E. Conrad and W.C. Oechel, Technical Coordinators. Proceedings of the symposium on the dynamics and management of Mediterranean-type ecosystems. USDA Forest Service, General Technical Report PSW-58
- , J.H. Davis, J.P. Marshal, S.G. Torres, and B.J. Gonzales. 2009. Mining activity and habitat use by mountain sheep. *European Journal of Wildlife Research*, 55:183–191.
- Boyce, W.M. and R.N. Brown. 1991. Antigenic characterization of *Psoroptes* spp. (Acari: Psoroptidae) mites from different hosts. *Journal of Parasitology*, 77:675–679.
- Clark, R.K., D.A. Jessup, and R.A. Weaver. 1988. Scabies mite infestation in desert bighorn sheep from California. *Desert Bighorn Council Transactions*, 32:13–15.
- , W.M. Boyce, D.A. Jessup, and L.F. Elliott. 1993. Survey of pathogen exposure among population clusters of bighorn sheep (*Ovis canadensis*) in California. *Journal of Zoo and Wildlife Medicine*, 24: 48–53.
- Dougherty, M.J. and J.D. McInerney. 2009. What's the denominator? A lesson on risk. *American Biology Teacher*, 71:31–33.
- Hanley, J.A. and A. Lippman-Hand. 1983. If nothing goes wrong, is everything all right? Interpreting zero numerators. *Journal of the American Medical Association*, 249:1743–1745.
- Ho, A.K. 2009. When the numerator is zero: another lesson on risk. *American Biology Teacher*, 71: 531–533.
- Hoban, P. 1990. A review of desert bighorn sheep in the San Andres Mountains, New Mexico. *Desert Bighorn Council Transactions*, 34:14–22.
- Jessup, D.A., W.E. Clark, and M.A. Fowler. 1986. *Wildlife restraint handbook*. Third edition. California Department of Fish and Game, Rancho Cordova, USA.

- , R.K. Clark, R.A. Weaver, and M.D. Kock. 1988. Safety and cost-effectiveness of net-gun capture of desert bighorn sheep. *Journal of Zoo and Wildlife Medicine*, 19:208–213.
- Lange, R.E., A.V. Sandoval, and S.P. Meleney. 1980. Psoroptic scabies in bighorn sheep (*Ovis canadensis mexicana*) in New Mexico. *Journal of Wildlife Diseases*, 16:77–82.
- Logan, K.A. and L.L. Swenor. 2001. Desert puma: evolutionary ecology and conservation of an enduring carnivore. Island Press, Covelo, California, USA. 463 pp.
- Mazet, J.A., W.M. Boyce, J. Mellies, I.A. Gardner, R.K. Clark, and D.A. Jessup. 1992. Exposure to *Psoroptes* sp. mites is common among bighorn sheep (*Ovis canadensis*) populations in California. *Journal of Wildlife Diseases*, 28:542–547.
- Monteith, K.M., V.C. Bleich, T.R. Stephenson, B.M. Pierce, M.M. Conner, R.W. Klaver, and R.T. Bowyer. 2011. Timing of seasonal migration in mule deer: effects of climate, plant phenology, and life-history characteristics. *Ecosphere*, 2(4):art47. DOI:10.1890/ES10-00096.1
- , T.R. Stephenson, V.C. Bleich, M.M. Conner, B.M. Pierce, and R.T. Bowyer. 2013. Risk-sensitive allocation in seasonal dynamics of fat and protein reserves in a long-lived mammal. *Journal of Animal Ecology*, 82:377–388.
- Roug, A., P. Swift, S. Torres, K. Jones, and C.K. Johnson. 2012. Serosurveillance for livestock pathogens in free-ranging mule deer (*Odocoileus hemionus*). Accessed 29 December 2013. PLoSONE, 7(11): e50600. doi:10.1371/journal.pone.0050600
- Schaefer, R.J., S.G. Torres, and V.C. Bleich. 2000. Survivorship and cause-specific mortality in sympatric populations of mountain sheep and mule deer. *California Fish and Game*, 86:127–135.
- Singer, R.S., D.A. Jessup, I.A. Gardner, and W.M. Boyce. 1997. Pathogen exposure patterns among sympatric populations of bighorn sheep, mule deer, and cattle. *Journal of Wildlife Diseases*, 33: 377–382.
- Wehausen, J.D. 1987. Some probabilities associated with sampling for diseases in bighorn sheep. *Desert Bighorn Council Transactions*, 31:8–10.
- Welsh, G.W. and T.D. Bunch. 1983. Psoroptic scabies in desert bighorn sheep (*Ovis canadensis nelsoni*) from northwestern Arizona. *Journal of Wildlife Diseases*, 19:342–344.
- Zahler, M., W.M.L. Hendriks, A. Essing, H. Rinder, and R. Gothe. 2000. Species of the genus *Psoroptes* (Acari: Psoroptidae): a taxonomic consideration. *Experimental and Applied Acarology*, 24: 213–225.